## Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

## Listing of Claims:

Please cancel claims 1-183 without prejudice and add the following new claims.

- 184. (New) A cigarette comprising a portion of an increased-nicotine transgenic *Nicotiana tabacum* plant, wherein said cigarette is characterized by (i) a tar-to-nicotine yield ratio of between about 3 and about 8, as measured by the FTC or ISO method, and (ii) cigarette smoke having a pH of about 6.5 or lower.
- 185. (New) A cigarette according to claim 184, wherein said plant expresses at least one heterologous nucleic acid that up-regulates the production of nicotine in said transgenic plant or plant portion.
- 186. (New) A cigarette according to claim 185, wherein said plant expresses a heterologous nucleic acid encoding at least a segment of QPT and PMT.
- 187. (New) A method of producing a nicotine buffer, comprising:
- a) crossing an increased-nicotine transgenic *Nicotiana tabacum* plant with an increased-sugar *Nicotiana tabacum* plant;
- b) selecting an increased-nicotine and increased-sugar *Nicotiana tabacum* progeny plant; and
- c) producing a cigarette comprising a portion of said increased-nicotine and increased-sugar *Nicotiana tabacum* plant.
- 188. (New) The method according to claim 187, wherein said increased-nicotine transgenic *Nicotiana tabacum* plant is produced by:
- a) transforming a *Nicotiana tabacum* plant with a construct comprising, in the 5' to 3' direction, a promoter operably linked to a heterologous nucleic acid encoding an enzyme or transcription factor that upregulates nicotine synthesis;
  - b) regenerating transgenic Nicotiana tabacum plants from the transformed plant; and

- c) selecting a transgenic *Nicotiana tabacum* plant having increased-nicotine content relative to a non-transformed control plant.
- 189. (New) A method for producing a nicotine buffer, comprising
- a) providing a transgenic *Nicotiana tabacum* plant that expresses a heterologous nucleic acid sequence that confers elevated nicotine and fatty acid levels, relative to a non-transformed control plant, and
  - b) producing a cigarette that comprises a portion of said plant.
- 190. (New) The method according to claim 189, wherein step (a) comprises:
- 1) transforming a *Nicotiana tabacum* plant with (i) a first construct comprising, in the 5' to 3' direction, a promoter operably linked to a heterologous nucleic acid encoding an enzyme or transcription factor that upregulates nicotine synthesis; and (ii) a second construct comprising, in the 5' to 3' direction, a promoter operably linked to a heterologous nucleic acid encoding an enzyme or transcription factor that up-regulates fatty acid synthesis
  - 2) regenerating transgenic Nicotiana tabacum plants from the transformed plant; and
- 3) selecting a transgenic *Nicotiana tabacum* plant having increased-nicotine content and increased fatty acids relative to a non-transformed control plant.
- 191. (New) A method according to claim 189, wherein step (a) comprises:
- a) crossing an increased-nicotine transgenic *Nicotiana tabacum* plant with an increased fatty acid *Nicotiana* transgenic plant;
- b) selecting an increased-nicotine and increased fatty acid *Nicotiana tabacum* progeny plant; and
  - c) producing a cigarette comprising a portion of said progeny plant.
- 192. (New) A method for increasing nicotine and cured reducing sugar content in a *Nicotiana tabacum* plant, comprising:
- a) transforming a *Nicotiana* plant having elevated cured reducing sugar content with a nucleic acid conferring an increased-nicotine phenotype;

- b) regenerating transgenic Nicotiana tabacum plants from the transformed plant; and
- c) selecting a progeny plant having increased cured reducing sugar content and increased-nicotine content relative to a non-transformed control plant.
- 193. (New) A method of increasing alkaloid biosynthesis in a *Nicotiana tabacum* plant or part thereof, comprising:
- a) transforming a *Nicotiana tabacum* plant with at least one nucleic acid sequence, wherein said nucleic acid increases quinolate phosphoribosyl transferase and putrescine N-methyltransferase expression;
  - b) regenerating transgenic Nicotiana tabacum plants from the transformed plant; and
- c) selecting a plant having increased alkaloid biosynthesis, wherein said plant has increased quinolate phosphoribosyl transferase and putrescine N-methyltransferase expression relative to a non-transformed control plant.
- 194. (New) A tobacco plant characterized by increased nicotine levels and high cured reducing sugar content, wherein at least one transgene confers the increased nicotine levels.
- 195. (New) A tobacco plant according to claim 194, wherein said plant has a nicotine content greater than 3.5 percent and a cured reducing sugar content from between about 12 percent and about 30 percent.
- 196. (New) A method of making a tobacco product having lower levels of tobacco-specific nitrosamines, comprising:
- a) providing a reduced nicotine transgenic *Nicotiana tabacum* plant or plant portion, wherein said plant or plant portion expresses at least one heterologous nucleic acid that down-regulates nicotine production;
  - b) producing tobacco from said transgenic plant or plant portion;
  - c) adding nicotine to said tobacco; and
- d) producing a product comprising said tobacco, wherein said product has a lower level of tobaccospecific nitrosamines relative to a control product comprising a non-transformed control plant or plant portion.
- 197. (New) A product produced by the method of claim 196, wherein said product is cigarette tobacco.

- 198. (New) The tobacco product of claim 197, wherein said cigarette tobacco comprises filler having a level of tobacco-specific nitrosamines below about 0.5 microgram per gram of tobacco.
- 199. (New) The tobacco product of claim 198, wherein said level of tobacco-specific nitrosamines is less than 0.05 microgram (50 ppb) per gram of tobacco.
- 200. (New) A method of making reconstituted tobacco, comprising:
  - a) providing plant material selected from a group consisting of
  - (i) an increased-nicotine transgenic *Nicotiana tabacum* plant or plant portion, wherein said plant or plant portion expresses at least one heterologous nucleic acid that up-regulates the production of nicotine in said transgenic plant or plant portion;
  - (ii) a reduced-nicotine transgenic *Nicotiana tabacum* plant or plant portion, wherein said plant or plant portion expresses at least one heterologous nucleic acid that down-regulates the production of nicotine in said transgenic plant or plant portion;
    - (iii) deproteinized tobacco fiber or tobacco fiber portion; and
    - (iv) freeze-dried tobacco or tobacco portion; and
  - b) combining and reconstituting any combination of said plant material.
  - 201. (New) A cigarette comprising the reconstituted tobacco of claim 200, wherein said cigarette has a reduced yield of compounds selected from a group consisting of polycyclic aromatic hydrocarbons, tobaccospecific nitrosamines, benzo(a)pyrene, phenols, and catechols relative to a cigarette made from a control plant.